



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### Testosterone, facial symmetry and cooperation in the Prisoners' Dilemma

**Citation for published version:**

Sanchez-Pages, S & Turiegano, E 2010, 'Testosterone, facial symmetry and cooperation in the Prisoners' Dilemma' *Physiology & Behavior*, vol. 99, no. 3, pp. 355-361. DOI: 10.1016/j.physbeh.2009.11.013

**Digital Object Identifier (DOI):**

[10.1016/j.physbeh.2009.11.013](https://doi.org/10.1016/j.physbeh.2009.11.013)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Peer reviewed version

**Published In:**

*Physiology & Behavior*

**Publisher Rights Statement:**

NOTICE: this is the author's version of a work that was accepted for publication in *Physiology & Behavior*. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in *Physiology & Behavior*, 99(3), 355-361 doi: 10.1016/j.physbeh.2009.11.013

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



Testosterone, Facial Symmetry and Cooperation in the Prisoners' Dilemma

Santiago Sanchez-Pages<sup>1</sup> and Enrique Turiegano<sup>2</sup>

<sup>1</sup>: *Edinburgh School of Economics, University of Edinburgh, EH8 9JT, UK.*

<sup>2</sup>: *Departamento de Biología, Universidad Autónoma de Madrid, 28049, Madrid, Spain.*

Correspondence: S. Sanchez-Pages, Edinburgh School of Economics. 31 Buccleuch  
Place, EH8 9JT, Edinburgh, United Kingdom. +44 131 651 3005 (e-mail:  
[ssanchez@staffmail.ed.ac.uk](mailto:ssanchez@staffmail.ed.ac.uk)).

NOTICE: this is the author's version of a work that was accepted for publication in Physiology and Behaviour. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in: Physiology & Behavior, 99(3), 355-361. doi: <http://dx.doi.org/10.1016/j.physbeh.2009.11.013>

**Abstract:**

Recent research has analyzed how individual characteristics, like the exposure to different hormones and symmetry, affect decision-making and strategic behaviour. The present article investigates the effect of symmetry, of exposure to testosterone (T) in utero and during puberty and of current T on cooperation in a Prisoners' Dilemma Game (PDG). T is a hormone with well known effect on males' behaviour, and that promotes activities that seek to increase reproductive success. Fluctuating Asymmetry (FA) reflects the ability of the organism to maintain a stable development and it is usually employed as a variable reflecting genetic quality (low FA values are thought to signal higher genetic quality). Our results show that subjects with intermediate levels of second to fourth digit ratio (a proxy of exposure to T in utero) and with high FA cooperate more often in the PDG. We also observe that the latter effect is due to the fact that FA has an impact on subjects' expectations about the behaviour of their counterpart in the game. These results reinforce the described link between markers related to genetic quality and cooperative behaviour. This possible linkage of individual condition and pro-social behaviour in humans clearly merits further attention.

*Keywords:* Testosterone, Cooperation, Prisoners' dilemma, Fluctuating asymmetry, Facial masculinity, 2D:4D.

## 1. Introduction

Testosterone (T) is a steroid hormone which determines in males of different species their development, their reproductive physiology and several behaviours [1-4]. In general, T affects males by promoting behaviours that seek to increase reproductive success, like an increased territorial aggressiveness [5,6], competitiveness [7] or a stronger status-seeking drive [8].

In mammals, T exerts organizational effects on the brain during foetal sexual differentiation [9] and during puberty [10]. These critical periods of exposure may affect adult male behaviour, in addition to the current level of T. Thus, in order to fully understand the influence of T in adult male behaviour, it is essential to take into account the exposure to the hormone during these critical periods. There are some measures in adult men that can proxy for T exposure in utero and during puberty. These are, respectively, the second to fourth digit ratio (2D:4D; the ratio between the length of index, or second digit, and ring finger, or fourth digit) and facial masculinity. Plenty of evidence indirectly suggests that 2D:4D negatively correlates with foetal T [11-13] and the existence or a significant negative association between 2D:4D and fetal testosterone/estradiol ratio (T/E ratio) [14]. On the other hand, many masculine facial features develop during puberty under the influence of T [15,16]. Most studies have found no correlation between these variables and the current T level, although some controversy remains. 2D:4D seems unrelated to current T in normal adults [17] and to facial metric measures of masculinity [18,19]. Sexually dimorphic facial traits are not associated either with current T [20], but there is some evidence of a link between perceived masculinity and current T [21,22] and 2D:4D [23].

There is also a wide body of literature linking these three variables with typically masculine features and behaviours. 2D:4D is a predictor of the degree of expression of

sexually dimorphic and other sex-hormone mediated traits, like visuo-spatial ability or left hand preference, and some behaviors like increased aggressiveness or competitiveness [24-27]. In men, circulating T has been linked to behaviours like acquisition of status, aggression, sensation-seeking or interest in sex [8,28-30]. Finally, the degree of masculinity has been shown to have an effect on male behaviour [20,31] but, above all, it has been described as a good predictor of male attractiveness [32-34]. Evidence shows that more attractive people behave differently [35-39]. However, it is important to bear in mind that masculinity is not the only factor determining male attractiveness [40-42].

Fluctuating asymmetry (FA) is a departure from symmetry in traits that are symmetrical at the population level [43]. It is considered to be the result of developmental instability. It thus reflects the ability of the organisms to maintain a stable development of their morphology and to overcome possible perturbations. This ability is thought to be affected by genetic and environmental factors. Many studies show a link between symmetry and genetic quality [44]. In humans, facial symmetry has been proposed as a cue for heritable fitness benefits [45,46], and it is widely considered as attractive [47,48]. Some studies have found that FA positively correlates with facial masculinity [49-51]. This suggests that both characteristics subtly indicate genetic quality. Still, many other studies find no correlation between them [52-54]. FA can be thus related to human behaviour in many different ways, since it is expected that the genetic quality of individuals has an impact on behaviour [55-58]. In general, symmetrical men (with low FA) tend to be less cooperative and more competitive. This behavior is believed to be due to their superior phenotypic quality, which increases their likelihood of winning conflicts and reduces their need to receive help from others [58].

In this study, we examine the effect of proxies for T exposure in utero and during puberty, of current levels of T and of FA on how human males play a one-shot symmetric Prisoner Dilemma Game (PDG). The effects of some of these variables have been previously tested in different economic experiments, like the ultimatum game [58-60], public good games [61], the dictator game [62] and risk-taking in an investment game [31]. Formally, the symmetric PDG is a special case of the public good game with two players and two available actions: “Cooperate” (equivalent to a full contribution) and “Defect” (equivalent to no contribution). The outcome obtained by the players when both defect is worse for each of them than the outcome they would have obtained if both of them had cooperated. When the players choose different actions, the one who cooperated receives a very low payoff, while the defector obtains a very high payoff. “Defect” is thus a dominant strategy in this game, that is, it is the best strategy for both participants regardless of whatever their opponent does. However, it is well known that humans tend not to follow this rule. There is substantial experimental evidence showing that humans are willing to cooperate and trust others in the one-shot PDG [63,64].

We expect high FA males to cooperate more in the PDG, in line with some previous results [38,58,60]. As FA is a marker of genetic quality, high symmetric people have less need for receiving help from others, reducing their interest in mutual reciprocity. In the same line, as high T is also considered to be a marker of genetic quality, we expect the three T-related variables (2:4 finger ratio, Facial Masculinity and current T level) to have a positive impact on defection rates. Effects in this direction have been previously observed in other games [61]. However, it has been pro-social behaviour has also been observed in people with low 2D:4D [61,62]. Because of this, it is difficult to predict specifically what will be the sign of the effect on the PDG of the

exposure to high T at these three different stages [31], although it is likely that 2D:4D will display some effect [61].

## **2. Methods**

### *2.1. Participants*

The experiment was carried out in Edinburgh and Madrid. 160 students participated in the experimental sessions. We discarded answers from non-white students because we calculated facial masculinity by comparing each student photograph with an average image obtained from 50 photos of white female students (the most common racial group in our subject pool). In total, we employed 147 self-reported white students, 78 in Spain and 69 in Scotland. They were aged from 17 to 30, with the Spanish students ( $21.04 \pm 2.45$ ; mean  $\pm$  SD) being significantly older ( $t_{145} = 4.534$ ,  $p < 0.001$ ) than the Scottish ( $19.52 \pm 1.39$ ). Based on self reports, 139 subjects were heterosexuals and 8 were homosexual. Written consent was obtained from all participants and the collection of photographs, hand-scans and saliva was approved by the relevant ethics committees at each institution.

### *2.2. Experimental procedure*

The experiments were performed employing the z-Tree 3.2.10 software for Economics Experiments [65]. The experiments were run in sessions with less than 20 subjects. In order to avoid unexpected effects on participants' behaviour [59] all the experiments were tracked by the (male) authors. Before each session, all subjects were carefully instructed about the experiment and their photographs, hand-scans and saliva samples were taken. All the subjects filled a questionnaire asking their age, ethnicity, sexual orientation and degree. This study was part of a larger one that included several

other items. Apart from the show-up fee (£5 in Edinburgh, 5€ in Madrid), subjects were told that their final payment would depend upon their choices in several but not all of the items in the questionnaire. They were informed of which ones counted for payment only after the experiment concluded. Each experimental session took about an hour. Subjects were paid privately in cash after the session and after they filled the corresponding official receipt.

The PDG that subjects played was a one-shot game with two available strategies, “Cooperate” and “Defect”. If the two players choose “Cooperate” they both get 90 points, if both defect they both get 30 points. If they choose different actions, the one who cooperates gets 10 points and the one who defects obtains 160 points. Hence both players choosing “Defect” constitutes the unique Nash Equilibrium of the PDG. This strategy profile is also a Dominant Strategy Equilibrium, since “Defect” is a dominant strategy for both players. Subjects were asked which strategy they believed that their hypothetical counterpart would choose and also which action they would take.

### *2.3. Masculinity and FA Measurement*

Full frontal facial colour photographs were taken of all participants with an Olympus E-500 digital camera with resolution 3264x2448 in JPEG format under standardized light conditions. The camera distance was kept constant at 3m and the zoom was completely opened to avoid slight optical distortion of true facial shape. Participants were asked to remove any facial adornment, to pose with a neutral expression and to look directly into the camera. We took three images of each participant in order to choose the best one for our purposes. Facial measures, as masculinity or FA, were calculated from the photographs using geometric morphometric tools [66]. The shape of each face was defined by manually setting 39



predetermined points called landmarks (LM). These 39 points (Figure 1) were chosen because they can be unambiguously identified in every photo. This ensures that they mark positions which rigorously correspond, in a biological or perceptual ground, to the same position in every face [67]. The LMs were placed twice, once by each author, which makes possible to assess any measurement error.

We employed these LMs to calculate the Procrustes distance between pairs of rotated and scaled images [67] using the TPS software package (by F.J. Rohlf, see <http://life.bio.sunysb.edu/morph/>). To calculate the asymmetry of each image, we compared the LM position of each face and a mirror-image of the same one, measuring the Procrustes distance between both LM positions [68]. FA can be understood as a deviation of the “perfect” symmetry or, as it is commonly considered, as an individual deviation from the average (directional) asymmetry. In this context the asymmetry of a bilateral object is attributable partially to directional asymmetry (differences in the population between average right and left size) and partially to fluctuating asymmetry (deviation of each individual’s asymmetry from the overall average asymmetry). We obtained FA by decomposing the Procrustes distance between each image and its mirror reflecting in directional and fluctuating asymmetry by employing the Procrustes ANOVA method. The latter method characterizes the shape of an object (the faces) as a single geometric object. Because calculation of Procrustes coordinates is based on the algebra of sums of squares, individual deviations from the average shape can be partitioned in different components, as happens in the conventional ANOVA [69]. The classic ideas of fluctuating and directional asymmetry are applied using this alternative approach [70], where directional asymmetry corresponds to the variation introduced by the variable “side of the object”, while FA corresponds to the variation explained by the interaction between side and individual. To compute the FA of each individual we

employed Morpho J software (by C. P. Klingenberg. See [http://www.flywings.org.uk/MorphoJ\\_page.htm](http://www.flywings.org.uk/MorphoJ_page.htm)). The FA values obtained correlate strongly with the total asymmetry calculated for each face ( $r=0.982$ ,  $p<0.001$ ).

The masculinity of the faces was measured calculating the Procrustes distance between the LMs of the male half-faces (where only 22 LM keep placed) and the LMs of a female half-face obtained by averaging 50 images of female students. With this protocol, each male presented two masculinity distances, one for each hemi-face. The measure of masculinity employed in the analyses is the average of these two distances. We employed hemi-faces to calculate the masculinity in order to avoid incorporating the measure of the symmetry of the face indirectly (given that the female average image is completely symmetrical). To perform this protocol we randomly chose one of the two possible sets of LMs (one placed by each author). Both LMs configurations are strongly correlated ( $r=0.998$ ,  $p<0.001$ ). Masculinity understood as the difference in shape between standard male and female faces has been widely employed in order to generate feminized and masculinized faces [71,72].

#### 2.4. Digit Ratio Measurement

Participants' right hands were scanned with an Hp psc 2110 scanner with a resolution of 600x1200 ppi. The second and fourth digits were measured from the centre of the flexion crease proximal to the palm to the top of the digit. This is a commonly accepted way to calculate 2D:4D [31,61,67]. To measure the fingers both authors independently placed a LM in each of the described positions and both lengths were measured afterwards. The placing of LMs and the measures were done with the appropriate utility of the TPS morphometric free software package. The two

measurements of 2D:4D were highly correlated ( $r = 0.96$ ,  $p < 0.001$ ,  $N = 147$ ). The measure employed in the analysis was their average. In some cases, it was necessary to repeat the hand scanning because the image was unsuitable for correct measuring. We measured lengths in pixels up to two decimal places.

### 2.5. Salivary T Measurement

Current T was measured from saliva provided only by the subjects in Madrid, following the protocol suggested by previous studies [73]. Saliva samples were taken from each participant 30 minutes upon arrival in order to be sure that they have not eaten, drunk or brushed their teeth just before saliva sampling. All samples were collected between 11:00 and 13:00, and participants were asked to spit through a straw into a saliva sampling device (SALI-TUBES 100, DRG). No significant differences in T concentrations were found between subjects as a function of the hour in which the samples were collected. Saliva samples were immediately centrifuged, frozen and stored at  $-20^{\circ}\text{C}$ . At the end of the collection period, all samples were assayed employing T assays commercially available kits (Salivary T ELISA kit from DRG Diagnostics). Two kits were employed successively, and the sample concentrations used in the analyses are the averages of the duplicates. Inter-assay coefficients of variation were 14.26% and the intra-assay coefficient of variation was 10.87%. One of the saliva samples was discarded because it presented visible blood contamination. We were unable to obtain measures from two other subjects because there was not enough volume of sample to duplicate the measure.

### 2.6. Statistics

We tested the normality of all our variables with the Kolmogorov-Smirnov test. Salivary T and 2D:4D are normally distributed, but we had to log transform FA and masculinity after multiplying them by 100 (in order to avoid negative values that could interfere with the interpretation of their effects). To analyze the results we employed two-tailed Student-t tests. We also employed logistic regressions to analyze the effect of several independent variables on our dichotomous dependent variable (“Cooperate” or “Defect”). We employed SPSS12 for all the statistical analyses.

### 3. Results

Table 1 presents descriptive statistics for each variable. The t tests show that participants in each city do not differ in 2D:4D ( $t_{145}=0.263$ ,  $p=0.793$ ), FA ( $t_{145}=0.657$ ,  $p=0.512$ ) or facial masculinity ( $t_{145}=1.426$ ,  $p=0.156$ ). There exist age differences between both groups (see methods). We have found that Spanish subjects cooperate more often (62.82%) than the Scottish (42.03%;  $\chi^2_1=6.355$ ,  $p=0.012$ ). City was therefore used as a control variable in all further analyses.

No correlation was found between any combination of the three variables (FA, masculinity and 2D:4D) and age, except for a significant correlation between masculinity and FA ( $r=0.320$ ,  $p<0.001$ ).

Table 1 provides average measures of participants depending on whether they chose “Cooperate” or not, and the significance of the differences across these two groups. Participants who cooperated had significantly higher FA values than those who did not.

In order to simultaneously evaluate the effect of all the variables on cooperative behaviour we built a logistic regression model including City as a control variable, FA, and Masculinity. We also included 2D:4D and its second order term, in order to

correctly account for the non-linear effect of this variable [61]. The resulting model was significant (see Table 2). We found a highly significant effect of FA on cooperation. That is, men with higher FA levels tend to cooperate more in the PDG. We also found a significant effect of 2D:4D and its second term, positive and negative respectively, implying that men with intermediate values of 2D:4D are more likely to cooperate. Moreover, the model including solely 2D:4D and its second order term was also significant (see Table 2). These both effects can be roughly observed if we divided the sample into blocks. When we divide the sample in two equal-sized blocks according to FA, participants who presented high values of FA cooperated more often (61.12%) than low FA participants (45.34%). On the other hand, if the sample is divided in three equal-sized blocks according to 2D:4D, it is possible to observe that participants who showed an intermediate value of 2D:4D (the intermediate third of them) tend to cooperate more often (67.35%), whereas participants with low or high values cooperate less frequently (45.10% and 46.81% respectively).

As FA and masculinity correlates, we built a model excluding masculinity given that we found no differences in masculinity between those participants who cooperated and those who did not (see Table 2). In order to account for possible interactions between the variables (FA, 2D:4D, masculinity), we run several models including interacting terms but none of them were significant (not shown).

Another variable that can affect cooperation is the expected behaviour of the counterpart (EB). This variable is strongly significant (see Table 2) and its inclusion in our model renders City and FA insignificant, implying that these two variables are somehow related to EB. The participants who thought that the other part will cooperate

show higher FA than the rest ( $t_{145}=2.011$ ,  $p=0.046$ ) while no differences were found in City ( $\chi^2_1=3.183$ ,  $p=0.074$ ).

Salivary T levels were only measured for the Spanish subjects, and not for all of them ( $n=75$ ). Salivary T did not correlate with 2D:4D ( $r=-0.146$ ,  $p=0.210$ ), facial masculinity ( $r=0.069$ ,  $p=0.555$ ) nor FA ( $r=-0.087$ ,  $p=0.465$ ). We found no differences in Salivary T levels between participants who cooperated and those who did not (see Table 1). A model including only these 75 participants displays exactly the same features as the model that included all the participants, that is, the positive effect of FA on cooperation and that subjects with intermediate 2D:4D values tend to be more cooperative (see Table 2). The model that includes Salivary T is also statistically significant, but not the variable itself.

#### 4. Discussion

The objective of this study is to analyze the relationship between cooperative behaviour in the PDG and a set of individual characteristics, some of them related to the exposure to T during life. Our results show a link between two of these characteristics, FA and 2D:4D, and cooperative behaviour. Participants who showed an intermediate value of 2D:4D tend to cooperate more often, while the participants with high FA also cooperate more. These results are in line with the results obtained in other studies [38,58-60]. We found no relationship between cooperation in the PDG and current (salivary) T nor facial masculinity (our proxy for T exposure during puberty).

No previous studies have attempted to explore the link between FA and cooperation, although the relevance of FA in other behaviours is well known [55-58]. On the other hand, very few studies have analyzed the effect of T on cooperation, although its effects on human behaviour have been extensively investigated

[8,24,28,31,74]. The closest contributions to ours have studied the effect of these two variables on the Ultimatum Game [58-60,75]. The Ultimatum Game (UG) is not normally considered as a game of cooperation because it does not contain a fundamental tension between social and private incentives. Still, some authors have used it as an approximation to the cooperative interactions that occur during hunting [76]. Under this interpretation, a dominant individual tries to obtain the cooperation of another one in order to hunt, and proposes a division of the expected catch. The non-dominant individual can accept or reject that proposal. Rejection means that no catch is obtained. Note that, contrary to what happens in the PDG, social and private incentives are aligned when the second individual has to make a choice.

In the UG, males with low 2D:4D (presumably exposed to high T/E ratio in utero) have higher minimum acceptable offers, although there is no described relation between this variable and the offers made [59,60]. On the other hand, males with low FA (that is, more symmetric) make lower offers, although there is no described relation between this variable and the likelihood of rejection [58]. Our results are in line with these studies, as FA affects males' behaviour in the PDG by influencing the estimation that players have about the choice of the other player, similarly to what occurs in the UG (where symmetry affects offers, which in turn are an indirect measure of the expected probability of acceptance). Symmetric males thus tend to believe that their counterpart will defect and behave accordingly. This is in line with previous studies that have observed that symmetric males tend to cooperate less frequently because of their superior phenotypic quality [58]. Given their higher ability in obtaining resources, males with low FA do not need to be, nor look as cooperative as males with high FA, and then cooperate less. This is however at odds with other experiments in which more attractive subjects trust others more in a Trust Game [39]. This can be explained if we

bear in mind that attractiveness does not only depend on FA [34] and that in that experiment the subject pool was composed by males and females, and they tend to evaluated attractiveness attending to different features [77]. In any case the PDG and the Trust Game are very different games. In the Trust Game, as the second players acts after the first have trusted (or not), “leading by example” can have an important effect. This is not possible in the PDG because it is simultaneous. Symmetric males may then trust because of their self-confidence and because they may attempt to obtain higher status by leading cooperation.

We also find that individuals with lower 2D:4D, (exposed to higher T level in utero) tend to defect more, in the same line as in the UG [59,60]. This does not coincide completely with the results obtained in studies that explored the relationship between individual characteristics and contributions in a public good game [61]. They find that individuals which did not cooperate or behaved altruistically (that is, they contributed more than what the social norm dictates) show high 2D:4D, although these effects were not statistically significant in men. In our experiment, cooperation is just a dichotomous variable while in the public good game cooperation is a spectrum (i.e., the amount of the contribution). Our experimental design is thus simpler and can offer cleaner results but this comes at the price of limiting the richness of possible behaviours. We find, as these authors do, that low 2D:4D subjects are less likely to behave altruistically (tend to cooperate less often in our game). However, their conjecture that subjects with low 2D:4D tend to adhere to social norms cannot explain our results, since defecting is unlikely to be the social norm in PDG. In our case, self-sufficiency in obtaining resources can explain the behaviour of subjects with low 2D:4D. This point of view is in line with another study on cooperation in which more attractive males tended to cooperate less [38]. As we have already mentioned, we cannot assume that our more



symmetric or masculine subjects will be identified as the more attractive because this is a trait affected by many other variables [34]. In any case, our results go in the same direction as those.

Another interesting result is that individuals with high 2D:4D, that is, those less exposed to T in utero, tend to cooperate less often. This result has not been observed in any previous analysis and it is thus difficult explain within the domain of usual explanations. Unlike FA, 2D:4D has no impact on the behaviour that players expect from their counterparts. Hence, the lack of cooperation of these subjects cannot be attributed to their beliefs. This is an interesting result that we plan to explore properly in our future research. The significant differences in cooperation rates between Edinburgh and Madrid show that behaviour in PDG, as most human behaviours, is strongly affected by cultural constraints. But the biological features also have an important effect. Biological individual characteristics remain strongly significant after we control for cultural differences in our logistic model, and also when we restrict the analysis to only the Spanish subjects (see Table 2).

In our study, two other variables that could potentially affect cooperative behaviour, like facial masculinity and current T, display no effect. It is well known that the current level of T is linked to aggressiveness and status-seeking behaviour [8,28]. In addition, facial masculinity might have shown an effect as it is considered as a signal of genetic fitness according to the “immunocompetence handicap hypothesis” [79]. We have found no relationship between these variables and cooperation in the PDG. Hence, based on these lacks of effect we conclude that cooperation in PDG is not understood as a challenge [29,80] and that the aversion to a possible breakdown of cooperation cannot be equated to the standard concept of risk aversion [31]. It is interesting that the exposure to T in some periods of life seems to have an impact on certain types of

behaviour, like risk aversion [31], but not in others, like the ones we investigate in this study. This suggests that exposure to T influences behaviour in very diverse ways. The different effects of the levels of T during development and the links between the behaviours that seem to be affected by the hormone also deserve further experiments. To perform such studies it would be necessary to test the same pool of subjects in different economic experiments after controlling for all these variables. In the same line, to ensure that developmental instability is behind facial FA, it will be necessary complement further data on cooperation and FA with measures in some others bilateral traits. In addition, it would be necessary to extend the experiments by employing women and non-students as subjects, enlarging thus the range of age and occupations.

#### **ACKNOWLEDGMENTS**

The authors acknowledge the financial support of the Abbey/Santander Research Fund and thank I. Monedero, M. Pita and M. Losada for their help with the experiments, J. Marugan and C.P. Klingenberg for their help with Geometric Morphometry and FA measures and to R. Forshaw for her constructive comments on the paper.

## 5. References

- [1] Wingfield JC, Hegner RE, Dufty J, Ball GF. The "challenge hypothesis": theoretical implications for patterns of T secretion, mating systems, and breeding strategies. *Am Nat* 1990;136:829-846.
- [2] Oliveira RF, Ros AFH, Gonçalves DM. Intra-sexual variation in male reproduction in teleost fish: A comparative approach. *Horm Behav* 2005;48:430-439.
- [3] Woolley SC, Sakata JT, Crews D. Evolutionary insights into the regulation of courtship behavior in male amphibians and reptiles. *Physiol Behav* 2004;83:347-360.
- [4] Fusani L. Testosterone control of male courtship in birds. *Horm Behav* 2008;54:227-233.
- [5] Wingfield JC. A continuing saga: the role of Testosterone in aggression. *Horm Behav* 2005;48:253-255; discussion 256-258.
- [6] Soma KK, Scotti ML, Newman AEM, Charlier TD, Demas GE. Novel mechanisms for neuroendocrine regulation of aggression. *Front Neuroendocrinol* 2008;29:476-489.
- [7] Edwards DA. Competition and Testosterone. *Horm Behav* 2006;50: 681-683.
- [8] Mazur A, Booth A. Testosterone and dominance in men. *Behav Brain Sci* 1998;21:353-363.
- [9] Morris JA, Jordan CL, Breedlove SM. Sexual differentiation of the vertebrate nervous system. *Nat Neurosci* 2004;7:1034-1039.
- [10] Sisk CL, Schulz K, Zehr J. Puberty: A finishing school for male social behavior. *Ann NY Acad Sci* 2003;1007:189-198.

- 419 [11] Manning J, Scutt D, Wilson J, Lewis-Jones D. The ratio of 2nd to 4th digit length:  
420 a predictor of sperm numbers and concentrations of testosterone, luteinizing  
421 hormone and oestrogen. *Hum Reprod* 1998;13:3000-3004.
- 422 [12] Cohen-Bendahan CC, van de Beek C, Berenbaum SA. Prenatal sex hormone  
423 effects on child and adult sex-typed behavior: methods and findings. *Neurosci*  
424 *Biobehav R* 2005;29:353-384.
- 425 [13] McIntyre M. The use of digit ratios as markers for perinatal androgen action.  
426 *Reprod Biol Endocrin* 2006;4:10.
- 427 [14] Lutchmaya S, Baron-Cohen S, Raggatt P, Knickmeyer R, Manning JT. 2nd to 4th  
428 digit ratios, fetal testosterone and estradiol. *Early Hum Dev* 2004;77:23-28.
- 429 [15] Enlow D, Hans M. *Essential of facial growth*. Philadelphia: WB Saunders  
430 Company; 1996.
- 431 [16] Kasperk C, Helmboldt A, Börcsök I, Heuthe S, Cloos O, Niethard F, et al. Skeletal  
432 site-dependent expression of the androgen receptor in human osteoblastic cell  
433 populations. *Calcified Tissue Int.* 1997;61:464-473.
- 434 [17] Hönekopp J, Bartholdt L, Beier L, Liebert A. Second to fourth digit length  
435 ratio:2D:4D;and adult sex hormone levels: New data and a meta-analytic review.  
436 *Psychoneuroendocrinology* 2007;32:313-321.
- 437 [18] Koehler N, Simmons LW, Rhodes G. How well does second-to-fourth-digit ratio  
438 in hands correlate with other indications of masculinity in males? *Proc Biol Sci*  
439 2004; 271 Suppl 5:S296-S298.
- 440 [19] Burriss R, Little A, Nelson E. 2D:4D and sexually dimorphic facial characteristics.  
441 *Arch Sex Behav* 2007;36:377-384.
- 442 [20] Pound N, Penton-Voak IS, Surridge AK. Testosterone responses to competition in  
443 men are related to facial masculinity. *Proc Biol Sci* 2009;276:153-159.

- 444 [21] Penton-Voak IS, Chen JY. High salivary testosterone is linked to masculine male  
445 facial appearance in humans. *Evol Hum Behav* 2004;25:229-241.
- 446 [22] Roney JR, Hanson KN, Durante KM, Maestripieri D. Reading men's faces:  
447 women's mate attractiveness judgments track men's T and interest in infants. *Proc*  
448 *Biol Sci* 2006;273:2169–2175.
- 449 [23] Neave N, Laing S, Fink B, Manning JT. Second to fourth digit ratio, testosterone  
450 and perceived male dominance. *Proc Biol Sci* 2003;270:2167-2172.
- 451 [24] Manning J. Digit ratio: a pointer to fertility, behavior, and health. New Brunswick  
452 (NJ): Rutgers University Press; 2002.
- 453 [25] Putz DA, Gaulin SJC, Sporter RJ, McBurney DH. Sex hormones and finger length:  
454 What does 2D:4D indicate? *Evol Hum Behav* 2004;25:182-199.
- 455 [26] Hines M. Prenatal testosterone and gender-related behaviour. *Eur J Endocrinol*  
456 2006; 155:S115-S121.
- 457 [27] Hampson E, Ellis C, Tenk C. On the relation between 2D:4D and sex-dimorphic  
458 personality traits. *Arch Sex Behav* 2008;37:133-144.
- 459 [28] Roberti JW. A review of behavioral and biological correlates of sensation seeking.  
460 *J Res Pers* 2004;38:256-279.
- 461 [29] Archer J. T and human aggression: an evaluation of the challenge hypothesis.  
462 *Neurosci Biobehav Rev* 2006;30:319-345.
- 463 [30] Rupp HA, Wallen K. Relationship between testosterone and interest in sexual  
464 stimuli: The effect of experience. *Horm Behav* 2007;52:581-589.
- 465 [31] Apicella CL, Dreber A, Campbell B, Gray PB, Hoffman M, Little AC.  
466 Testosterone and financial risk preferences. *Evol Hum Behav* 2008;29:384-390.
- 467 [32] Thornhill R, Gangestad SW. Facial attractiveness. *Trends Cogn Sci* 1999;3:452-  
468 460.

- 469 [33] Johnston VS. Mate choice decisions: the role of facial beauty. *Trends Cogn Sci*  
470 2006;10:9-13.
- 471 [34] Rhodes G. The evolutionary psychology of facial beauty. *Annu Rev Psychol*  
472 2006;57:199-226.
- 473 [35] Mulford M, Orbell J, Shatto C, Stockard J. Physical attractiveness, opportunity,  
474 and success in everyday exchange. *Am J Sociol* 1998;103:1565-1592.
- 475 [36] Langlois JH, Kalakanis L, Rubenstein AJ, Larson A, Hallam M, Smoot M.  
476 Maxims or myths of beauty? A meta-analytic and theoretical review. *Psychol Bull*  
477 2000;126:390-423.
- 478 [37] Kanazawa S, Kovar JL. Why beautiful people are more intelligent. *Intelligence*  
479 2004;32:227-243.
- 480 [38] Takahashi C, Yamagishi T, Tanida S, Kiyonari T, Kanazawa S. Attractiveness and  
481 cooperation in social exchange. *Evol Psych* 2006;4:315-329.
- 482 [39] Wilson RK, Eckel CC. Judging a book by its cover: Beauty and expectations in the  
483 trust game. *Polit Res Q* 2006;59:189-202.
- 484 [40] Penton-Voak I, Jacobson A, Trivers R. Populational differences in attractiveness  
485 judgements of male and female faces: Comparing British and Jamaican samples.  
486 *Evol Hum Behav* 2004;25:355-370.
- 487 [41] Buss D. Strategies of human mating. *Psychol Top* 2006;15:239-260.
- 488 [42] Ellison PT. Beauty: In the gonads of the beholder - and the beheld. *Horm Behav*  
489 2008;53:11-13.
- 490 [43] Valen LV. A study on fluctuating asymmetry. *Evolution* 1962;16:125-142.
- 491 [44] Møller AP. Developmental stability and fitness: a review. *Am Nat* 1997;149:916-  
492 932.

- 493 [45] Scheib JE, Gangestad SW, Thornhill R. Facial attractiveness, symmetry and cues  
494 of good genes. *Proc Biol Sci* 1999;266:1913-1917.
- 495 [46] Little AC, Jones BC. Attraction independent of detection suggests special  
496 mechanisms for symmetry preferences in human face perception. *Proc Biol Sci*  
497 2006;273:3093-3099.
- 498 [47] Perrett DI, Burt DM, Penton-Voak IS, Lee KJ, Rowland DA, Edwards R.  
499 Symmetry and human facial attractiveness. *Evol Hum Behav* 1999;20:295-307.
- 500 [48] Little AC, Apicella CL, Marlowe FW. Preferences for symmetry in human faces in  
501 two cultures: data from the UK and the Hadza, an isolated group of hunter-  
502 gatherers. *Proc Biol Sci* 2007;274:3113-3117.
- 503 [49] Gangestad SW, Thornhill R. Facial masculinity and fluctuating asymmetry. *Evol*  
504 *Hum Behav* 2003;24:231-241.
- 505 [50] Brown WM, Price ME, Kang J, Pound N, Zhao Y, Yu H. Fluctuating asymmetry  
506 and preferences for sex-typical bodily characteristics. *Proc Nat Acad Sci*  
507 2008;105:12938-12943.
- 508 [51] Little AC, Jones BC, Waitt C, Tiddeman BP, Feinberg DR, Perrett DI, et al.  
509 Symmetry is related to sexual dimorphism in faces: data across culture and  
510 species. *PLoS ONE* 2008;3:e2106.
- 511 [52] Penton-Voak IS, Jones BC, Little AC, Baker S, Tiddeman B, Burt DM, et al.  
512 Symmetry, sexual dimorphism in facial proportions and male facial attractiveness.  
513 *Proc Biol Sci* 2001;268:1617-1623.
- 514 [53] Koehler N, Simmons LW, Rhodes G, Peters M. The relationship between sexual  
515 dimorphism in human faces and fluctuating asymmetry. *Proc Biol Sci* 2004;271  
516 Suppl 4:S233-236.

- 517 [54] Van Dongen S, Ten Broek CMA, Galis F, Wijnaendts LCD. No association  
518 between fluctuating asymmetry in highly stabilized traits and second to fourth digit  
519 ratio (2D:4D) in human fetuses. *Early Hum Dev* 2009;85:393-398.
- 520 [55] Furlow B, Gangestad SW, Armijo-Prewitt T. Developmental stability and human  
521 violence. *Proc Biol Sci* 1998;265:1-6.
- 522 [56] Manning J, Wood D. Fluctuating asymmetry and aggression in boys. *Hum Nat*  
523 1998;9:53-65.
- 524 [57] Pound N, Penton-Voak IS, Brown WM. Facial symmetry is positively associated  
525 with self-reported extraversion. *Pers Individ Differ* 2007;43:1572-1582.
- 526 [58] Zaatari D, Trivers R. Fluctuating asymmetry and behavior in the ultimatum game  
527 in Jamaica. *Evol Hum Behav* 2007;28:223-227.
- 528 [59] Van den Bergh B, Dewitte S. Digit ratio (2D:4D) moderates the impact of sexual  
529 cues on men's decisions in ultimatum games. *Proc Biol Sci* 2006;273:2091-2095.
- 530 [60] Burnham TC. High-T men reject low ultimatum game offers. *Proc Biol Sci*  
531 2007;274:2327-2330.
- 532 [61] Millet K, Dewitte S. Second to fourth digit ratio and cooperative behavior. *Biol*  
533 *Psychol* 2006;71:111-115.
- 534 [62] Millet K, Dewitte S. The presence of aggression cues inverts the relation between  
535 digit ratio (2D:4D) and prosocial behaviour in a dictator game. *Brit J Psychol*.  
536 2009;100:151-162.
- 537 [63] Marwell G, Ames RE. Economists free ride, does anyone else? : Experiments on  
538 the provision of public goods, IV. *J Public Econ* 1981;15:295-310.
- 539 [64] Dawes RM, Thaler RH. Anomalies: Cooperation. *J Econ Perspect* 1988;2:187-197.
- 540 [65] Fischbacher U. z-Tree: Zurich toolbox for ready-made economic experiments. *Exp*  
541 *Econ* 2007;10:171-178.



- 542 [66] Bookstein FL. Morphometric tools for landmark data: geometry and biology.  
543 Cambridge University Press; 1991.
- 544 [67] Fink B, Grammer K, Mitteroecker P, Gunz P, Schaefer K, Bookstein FL, et al.  
545 Second to fourth digit ratio and face shape. 2005;Proc Biol Sci 272:1995-2001.
- 546 [68] Klingenberg CP, Barluenga M, Meyer A. Shape analysis of symmetric structures:  
547 quantifying variation among individuals and asymmetry, Evolution 2002;56:1909-  
548 1920.
- 549 [69] Klingenberg CP, McIntyre GS. Geometric morphometrics of developmental  
550 instability: analyzing patterns of fluctuating asymmetry with procrustes methods.  
551 Evolution 1998;52:1363-1375.
- 552 [70] Schaefer K, Lauc T, Mitteroecker P, Gunz P, Bookstein FL. Dental arch  
553 asymmetry in an isolated Adriatic community. Am J Phys Anthropol  
554 2006;129:132-142.
- 555 [71] Perrett DI, Lee KJ, Penton-Voak I, Rowland D, Yoshikawa S, Burt DM, et al.  
556 Effects of sexual dimorphism on facial attractiveness. Nature 1998;394:884-887.
- 557 [72] Penton-Voak IS, Perrett DI, Castles DL, Kobayashi T, Burt DM, Murray LK, et al.  
558 Menstrual cycle alters face preference. Nature 1999;399:741-742.
- 559 [73] Granger DA, Shirtcliff EA, Booth A, Kivlighan KT, Schwartz EB. The "trouble"  
560 with salivary T. Psychoneuroendocrinology 2004;29:1229-1240.
- 561 [74] Coates JM, Herbert J. Endogenous steroids and financial risk taking on a London  
562 trading floor. Proc Nat Acad Sci 2008;105:6167-6172.
- 563 [75] Zaatari D, Palestis BG, Trivers R. Fluctuating asymmetry of responders affects  
564 offers in the ultimatum game oppositely according to attractiveness or need as  
565 perceived by proposers. Ethology 2009;115:627-632.

- 566 [76] Page KM, Nowak MA, Sigmund K. The spatial ultimatum game. *Proc Biol Sci*  
567 2000;267:2177-2182.
- 568 [77] Marcus DK, Miller RS. Sex differences in judgments of physical attractiveness: A  
569 social relations analysis. *Pers Soc Psychol Bull* 2003;29:325-335.
- 570 [78] Folstad I, Karter AJ. Parasites, bright males, and the immunocompetence  
571 handicap. *Am Nat* 1992;139:603.
- 572 [79] Mehta PH, Josephs RA. Testosterone change after losing predicts the decision to  
573 compete again. *Horm Behav* 2006;50:684-692.

Tables

	TOTAL <sup>a</sup>	Defect	Cooperate		
Age (yr) <sup>b</sup>	20.327±2.115	20.014±1.898	20.603±2.337	t <sub>145</sub> =-1.661	p=0.099
2:4 finger ratio <sup>b</sup>	0.962±0.030	0.965±0.036	0.959±0.025	t <sub>145</sub> =1.097	p=0.274
Fluctuating Asymmetry <sup>b</sup>	0.035±0.013	0.032±0.012	0.037±0.014	t <sub>145</sub> =-2.473	p=0.015
Facial masculinity <sup>b</sup>	0.098±0.022	0.098±0.024	0.098±0.020	t <sub>145</sub> =-0.055	p=0.956
Salivary T (pg/ml) <sup>c</sup>	135.997±26.124	132.322±28.057	138.186±24.954	t <sub>73</sub> =-0.940	p=0.351

Table 1.

*Mean values of variables in the total population and according to participants' choice*

<sup>a</sup> Data are the mean (±SD)

<sup>b</sup> In our whole population, 69 individuals defected and 78 cooperated.

<sup>c</sup> Salivary T was only measured in the experimental sessions performed in Madrid, where 28 individuals defected and 47 cooperated.

584

Variables in the model	MODEL				VARIABLE				
	-2LL	Likelihood Ratio Test	df	p	variables	coef	Wald	df	P
<i>2D:4D, (2D:4D)<sup>2</sup></i>	193.048	10.186	2	0.006	Constant	-414.551	7.308	1	0.007
					2D:4D	865.059	7.392	1	0.007
					(2D:4D) <sup>2</sup>	-450.728	7.466	1	0.006
<i>2D:4D, (2D:4D)<sup>2</sup>, FA, Masculinity, City</i>	179.952	23.282	5	<0.001	Constant	-441.869	7.101	1	0.008
					2D:4D	921.353	7.198	1	0.007
					(2D:4D) <sup>2</sup>	-480.065	7.287	1	0.007
					FA	1.464	7.048	1	0.008
					Masculinity	-0.806	0.892	1	0.345
<i>2D:4D, (2D:4D)<sup>2</sup>, FA, City</i>	180.858	22.376	4	<0.001	Constant	-447.523	165.366	1	0.007
					2D:4D	929.505	342.558	1	0.007
					(2D:4D) <sup>2</sup>	-484.184	177.413	1	0.006
					FA	1.317	0.528	1	0.013
					City	0.810	0.357	1	0.023
<i>2D:4D, (2D:4D)<sup>2</sup>, FA, City, EB</i>	120.438	82.796	5	<0.001	Constant	-557.024	7.007	1	0.008
					2D:4D	1165.062	7.122	1	0.008
					(2D:4D) <sup>2</sup>	-607.991	7.220	1	0.007
					FA	0.959	2.444	1	0.118
					City	0.672	2.111	1	0.146
<i>2D:4D, (2D:4D)<sup>2</sup>, FA</i>	90.766	12.179	3	0.007	EB	-3.307	41.259	1	<0.001
					Constant	-582.708	5.329	1	0.021
					2D:4D	1208.706	5.379	1	0.020
					(2D:4D) <sup>2</sup>	-627.815	5.446	1	0.020
<i>2D:4D, (2D:4D)<sup>2</sup>, FA, Current T</i>	86.483	12.623	4	0.013	FA	1.709	5.781	1	0.016
					Constant	-572.817	4.432	1	0.035
					2D:4D	1181.555	4.426	1	0.035
					(2D:4D) <sup>2</sup>	-612.624	4.459	1	0.035
					FA	1.860	6.514	1	0.011
					Current T	0.014	1.884	1	0.170

ALL PARTICIPANTS

MADRID

585

586 **Table 2.**587 *Logistic models for the whole sample and for the Spanish subjects.*

588 The variables included are 2D:4D (second to fourth finger ratio) FA (fluctuating

589 asymmetry), Masculinity (facial masculinity), Current T (current salivary T) and EB

590 (expected behaviour of the counterpart).

591 **Figures**

592

593 **Figure 1.**

594 *Landmarks placement .*

595 A) An average face with the 39 landmarks placed. B) All 147 landmarks configurations  
596 superimposed after Procrustes Fit. These coordinates are the basis for all FA  
597 calculations.

598